



Gastroenteropancreatic Neuroendocrine Neoplasms (GEP-NEN)

David B. Zhen, MD
Associate Professor of Medicine
GI Medical Oncology
Director, Neuroendocrine Tumor Program
Fred Hutch Cancer Center | University of Washington
October 6, 2025

Disclosures


- **Advisory Board/Consulting:** Exelixis, Boehringer Ingelheim, Bristol Myers Squibb, Legend Biotech
- **Independent Data Monitoring Committee:** Jazz Pharmaceuticals/Zymeworks, Inc
- **Research Funding:** AstraZeneca, Bayer, Bristol Myers Squibb, Clasp Therapeutics, Cornerstone Pharmaceuticals, Daiichi-Sankyo, Eli Lilly, Roche/Genentech, Legend Biotech, Merck, Pfizer

Outline

- Overview and Clinical Symptoms of GEP-NEN
- Workup and Staging of GEP-NEN
- Treatments for GEP-NET
- Treatments for GEP-NEC

Overview and Clinical Presentations of GEP-NEN

Neuroendocrine Neoplasms (NEN)



- Neuroendocrine cells found through various body sites
- Produce hormones and peptides with biological activity
- NEN arise in different organs
- GI tract and lung are common sites of origin for NEN
- Some cases of unknown primary
- GI NEN often referred as gastroenteropancreatic (GEP-NEN)


<https://my.clevelandclinic.org/health/diseases/22006-neuroendocrine-tumors-net>

2022 WHO Pathological Classification of GEP-NEN

Differentiation

Well differentiated
Neuroendocrine tumor (NET)

New category compared to
prior WHO classifications



Poorly Differentiated
Neuroendocrine carcinoma (NEC)

Proliferation Indices

Ki-67 <3%
Mitotic index <2/HPF

Ki-67 3 – 20%
Mitotic index <2-20/HPF

Ki-67 >20%
Mitotic index >20/HPF

Ki-67 >20%
Mitotic index >20/HPF

Designation

Low grade/
Grade 1

Intermediate grade/
Grade 2


High grade/
Grade 3

High grade by default

Subclassified by histology

- Small Cell
- Large Cell

Relevance of 2022 WHO GEP-NEN Pathological Criteria




Tang et al. Clin Cancer Res 2015; 22:1011.

WD: Well differentiated, PD: Poorly differentiated
Gr: Grade; HG: High grade


- Prognosis:
 - WD-Gr1/2 NET: Years (Median ~12 years)
 - PD-NEC: <12 months
 - WD-Gr 3 NET: In between the above
- WD-Gr3 NET mutational profiles more similar to WD-Gr1/2 NET
 - NET: MEN1, DAXX, ATRX
 - **NEC: TP53, RB1 (akin to SCLC, can be used to differentiate WD-Gr3 from NEC)**
- WD-Gr3 NET less responsive to platinum/etoposide compared to PD-NEC
- Differentiating WD-Gr3 from PD-NEC is important for prognostic and treatment considerations

Epidemiology of GEP-NET

A All NETs and malignant neoplasms



B NETs by site



Multiple factors likely contributing to increased incidence but most likely due to increased awareness/diagnosis

Hereditary Syndromes Associated with GEP-NET

- Most cases of GEP-NET don't have an obvious risk factor
- **~10-20% of pNET are associated with hereditary syndrome**
- Hereditary associated pNETs tend to more indolent
- **Importance of screening for other cancers in these patients!!**

Inherited disorders associated with pancreatic neuroendocrine tumors

Syndrome	Associated clinical features	Chromosomal location	Pancreatic neuroendocrine tumor type	Lifetime Risk
MEN1	Primary hyperparathyroidism Pituitary tumors Less commonly <ul style="list-style-type: none"> ▪ Adrenocortical tumors ▪ Carcinoid tumors ▪ Nonmedullary thyroid tumors 	11q13	Nonfunctional Gastrinoma Insulinoma Various	80-100%
Von Hippel-Lindau disease (VHL)	Pheochromocytoma (often bilateral) Retinal and cerebellar hemangioblastomas Renal cell carcinoma	3p25-26	Nonfunctional Various, including cystic tumors	~20%
Neurofibromatosis 1 (von Recklinghausen disease)	Neurofibromas Café au lait spots Pheochromocytoma	17q11.2		~10%
Tuberous sclerosis	Cardiac rhabdomyomas Renal cysts Angiomyolipomas	9q33.34 and 16p13.3		~1%

Symptoms of Hormonal Excess in GEP-NET

- GEP-NET may produce and secrete hormones & neuromodulators causing symptoms
- Classified as functional vs. non-functional
- Symptoms do not correlate with tumor burden
- Treatment of hormone excess: somatostatin analogue (SSA), except insulinoma
- **Based on prevalence below, many patients have non-functional tumors (ie often asymptomatic)**



NOTE: Carcinoid Syndrome is less common in pNETs (i.e. hormone excess is not always “Carcinoid Syndrome”)

Carcinoids (WD Gr1/2 midgut NET) (8-35% functional)	pNETs (10-40% functional)
<ul style="list-style-type: none"> - Carcinoid syndrome → flushing, diarrhea, bronchoconstriction, carcinoid heart dx - Due to excess serotonin, tachykinins, or histamine - Typically associated with midgut NETs and in the setting of liver metastases 	<ul style="list-style-type: none"> - *Insulin (insulinoma) → hypoglycemia - *Gastrin (gastrinoma) → peptic ulcer disease - Vasoactive intestinal peptide (VIPoma) → diarrhea, hypokalemia - Glucagon (glucagonoma) → flushing, diarrhea, hyperglycemia

1. Choti et al. J Clin Oncol 2012;30:suppl abstr 4126. 2. Soga et al. J Exp Clin Cancer Res 1999;18:133.
3. Oberg K. Semin Oncol 2010;37:594. 4. Halfdanarson et al. Ann Oncol 2008;19:1727.

Clinical Symptoms of GEP-NET: Carcinoid Heart Disease

- ~1/3 of carcinoid syndrome-related deaths
- 20-65% of pts with carcinoid syndrome develop valvular pathology
- High serotonin and tachykinin levels released by carcinoid tumor cells → valvulitis and fibroblast proliferation
- Plaque-like fibrous thickening involving classically right heart valves (ie tricuspid)
- Treatment of carcinoid syndrome and management of heart failure
- **Valvular replacement needed if severe**
- **Need to screen for this if clinical concern (e.g. murmur, cardiac symptoms) and conduct surveillance echocardiogram if found**



Clinical Symptoms of NET: Fibrosis/Desmoplastic Reaction



Daskalakis K et al. Br J Surg, 2017: 104(1).




Figure: Role of the Microenvironment in the Pathogenesis of Neuroendocrine Tumors (NETs)—NET cells mutually interact with their microenvironment, prompting angiogenesis through cytokine secretion; inhibiting T-cell function by T-regulatory cell (Treg) dysregulation; promoting infiltration of mast cells via Myc upregulation; and driving fibroblast activation, which in turn enhances NET cell proliferation. CTGF = connective tissue growth factor; HIF-1 α = hypoxia inducible factor alpha; IL = interleukin; TGF = transforming growth factor; TH1 = T helper type 1 cell; VEGF = vascular endothelial growth factor. Information from References 19,22,23, and 81.

Cives M & Strosberg J. Oncology (Williston Park) 2014; Sep 28(9): 749-56, 758

- Excess hormone production can lead to a fibrotic/desmoplastic reaction
- Tethers nearby bowel and place patients at risk of bowel obstruction
- Need to monitor closely for symptoms of bowel obstruction (e.g. pain, cramping, difficulty BM's, N/V, etc)
- **Palliative resection of primary tumor (even if metastatic disease) often considered to avoid future risk of bowel obstruction**

Workup and Staging of GEP-NEN

Workup of GI NET

- **Goals of workup**
 - Assess primary site and stage
 - Characterize aggressiveness (grade, differentiation) – need tissue
 - Establish functionality
- **Testing modalities**
 - Imaging
 - Multiphase CT scan or MRI
 - Somatostatin receptor-based imaging: ^{68}Ga Dotatate-PET scan or ^{64}Cu Dotatate-PET scan
 - Endoscopy
 - Biochemical evaluation as clinically indicated (if suspicious symptoms present)

Importance of Multiphase CT Imaging

© 2020 UpToDate, Inc. and/or its affiliates. All Rights Reserved.


CT of neuroendocrine tumor liver metastases

Arterial Phase

- ~20 sec post-contrast injection
- Most NET seen better on this phase

Portal venous phase

- ~70 sec post-contrast injection
- Better for adenocarcinoma & some NET



In patient A (top two images), the hypervascular liver metastases are more clearly observed on the arterial phase (left) compared to the portal venous phase (right); whereas in patient B (lower two images), the liver metastases are not as hypervascular and more clearly delineated on the portal venous phase (right) compared to the arterial phase (left).

Basics of Somatostatin Receptor (SSTR) Imaging





Table 1: Somatostatin Analogs Currently in Use for SSTR PET/CT


Compound	Abbreviation	Receptor Subtypes
^{68}Ga -DOTA-Tyr ³ -octreotate	^{68}Ga -DOTATATE (GaTate)	SSTR 2
^{68}Ga -DOTA-Nal ³ -octreotide	^{68}Ga -DOTANOC (GaNoc)	SSTR 3, SSTR 5
^{68}Ga -DOTA-TyI ³ -octreotide	^{68}Ga -DOTATOC (GaToc)	SSTR 5

Hofman MS, Lau WFE, Hicks RJ. RadioGraphics 2015; 35:500–516

Sideris L, Dube P, Rinke A. The Oncologist 2012; 17: 747-755

- NET often express SSTR on the surface of NET cells
- There are 5 isoforms of SSTR (ie SSTR 1-5); most relevant in NET SSTR 2 (i.e. dotaTATE)
- Take advantage of this by developing a radiolabel that bind to SSTR on NET cell surface

SSTR Imaging: ^{68}Ga Dotatate is Standard



- Old standard was OctreoScan ($^{111}\text{Indium}$ pentotretotide radiolabel)
- ^{68}Ga -DOTATATE has increased sensitivity and more convenient
- Combined with PET/CT scans allow for better imaging visualization
- ^{68}Ga -DOTATATE FDA approved 2016 and
- **Dotatate-PET scans are standard of care (and superior to OctreoScan)**

⁶⁴Cu-dotatate PET scan

- Limitations of ⁶⁸Ga dotatate
 - Short half-life (1.1 hr)
 - Needs to be locally produced via a generator
 - Thus needs to be given close to time of scan (Potentially limiting scan availability)
- ⁶⁴Cu dotatate has a longer half life (12.7 hr), eliminating need for a generator and increase scan availability
- Studies show same quality and safety as ⁶⁸Ga dotatate PET scans
- FDA approved in 9/2020
- While either ⁶⁸Ga dotatate and ⁶⁴Cu dotatate can be used interchangeably, consistency of use of one modality helps with radiology interpretations

Differentiation and Somatostatin Receptor Expression

- **WD-lower grade NET**

- +⁶⁸Ga-dotatate PET avid (express SSTR)
- Negative on FDG-PET

- **High grade NET/PD-NEC**

- Often neg ⁶⁸Ga-dotatate PET (little to no expression of SSTR)
- Positive on FDG-PET



- **In some instances, both FDG-PET scan and ⁶⁸Ga-dotatate PET scan can be helpful**

- Determine NET de-differentiation (ie to higher grade status)
- Guide treatment options



WD-G1 NET rectal primary seen only on dotatate-PET

Mixed FDG avid and dotatate avid lesions in liver (Mixed PD-NEC and WD-G2 NET)

F-18
FDG
PET



⁶⁸Ga
Dotatate
PET



*Images from patient with mixed WD-Gr1 NET of the rectum and mixed WD-NET and PD-NEC in liver

Biochemical Testing



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 1.2023
Neuroendocrine and Adrenal Tumors

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

PRINCIPLES OF BIOCHEMICAL TESTING¹⁻¹⁰

- Some NETs can secrete specific neuroendocrine hormones. Hormonal workup should be guided by the presence of symptoms of the excess hormone. Screening for hormones in individuals with asymptomatic disease is not routinely required.
- Patients with functional tumors have clinical symptoms related to tumor-associated hormone excess.
- PPIs, other drugs, some medical conditions, and certain foods are known to cause false elevations in serum gastrin and chromogranin A. To confirm diagnosis, serum gastrin should ideally be checked when fasting and off PPI for >1 week. However, PPI should be continued in patients with overt clinical symptoms of gastrinoma and/or risks of complications.
- If MEN2 is suspected, then patients should be evaluated for pheochromocytoma/paraganglioma prior to any procedures.⁹

Syndrome	Location	Clinical Signs or Symptoms	Testing
Carcinoid syndrome (NETs of Gastrointestinal Tract)	Primary tumors in small bowel and appendix; rarely in rectum	<ul style="list-style-type: none"> • Primary tumors in the GI tract usually are not associated with symptoms of hormone hypersecretion unless extensive metastasis. • Symptoms of hormone secretion may include flushing, diarrhea, cardiac valvular fibrosis, and bronchoconstriction 	<ul style="list-style-type: none"> • 24-hour urine or plasma 5-HIAA • Foods to avoid for 48 hours prior to and during testing: avocados, bananas, cantaloupe, eggplant, pineapples, plums, tomatoes, hickory nuts/pecans, plantains, kiwi, dates, grapefruit, honeydew, or walnuts
Carcinoid syndrome (NETs of Lung and Thymus)	Primary tumors in lung or thymus	<ul style="list-style-type: none"> • Bronchial/thymic tumors may be associated with classic carcinoid syndrome as well as hypercortisolemia (± Cushing's syndrome) 	<ul style="list-style-type: none"> • 24-hour urine or plasma 5-HIAA • Foods to avoid for 48 hours prior to and during testing: avocados, bananas, cantaloupe, eggplant, pineapples, plums, tomatoes, hickory nuts/pecans, plantains, kiwi, dates, grapefruit, honeydew, or walnuts • Test for hypercortisolemia (± Cushing's syndrome) (NE-C 2 of 4)
Insulinoma	Pancreas	Hypoglycemia	<ul style="list-style-type: none"> • While hypoglycemic: • Serum insulin • Pro-insulin • C-peptide • See Workup for insulinoma (PanNET-5)
VIPoma	Most common in pancreas, rarely extra pancreatic	Severe watery diarrhea, hypokalemia	Serum VIP
Glucagonoma	Pancreas	Flushing, diarrhea, hyperglycemia, dermatitis, hypercoagulable state	Serum glucagon
Gastrinoma	Pancreas or duodenum	Gastric ulcers, duodenal ulcers, diarrhea	Serum gastrin ^a
Somatostatinoma	Pancreas or duodenum	Hyperglycemia, cholelithiasis, diarrhea/steatorrhea	Serum somatostatin

[Footnotes on NE-C 3 of 4](#)

[References on NE-C 4 of 4](#)

NE-C
1 OF 4

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

Version 1.2023, 06/02/2023 © 2023 National Comprehensive Cancer Network® (NCCN®). All rights reserved. NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN.

- Generally workup is guided by symptoms
- For most asymptomatic patients, hormone levels don't change management
- Often use as biomarkers for response (e.g. Chromogranin A), but remember they can fluctuate due to other factors
- Some exceptions for asymptomatic patients (e.g. screening for Cushing's syndrome/pituitary tumors in setting of MEN1)
- **Assessment of response and treatment should be based on the entire clinical picture (not just biochemical testing)**

Updates to AJCC v9 Staging for GEP-NET

Critical updates in neuroendocrine tumors: Version 9
American Joint Committee on Cancer staging system for
gastroenteropancreatic neuroendocrine tumors

Aman Chauhan MD¹ | Kelley Chan MD² | Thorvardur R. Halfdanarson MD³ |
Andrew M. Bellizzi MD⁴ | Guido Rindi MD, PhD^{5,6} | Dermot O'Toole MD⁷ |
Phillip S. Ge MD⁸ | Dhanpat Jain MD⁹ | Arvind Dasari MD¹⁰ |
Daniel A. Anaya MD, MSHCT¹¹ | Emily Bergsland MD¹² | Erik Mittra MD, PhD¹³ |
Alice C. Wei MD¹⁴ | Thomas A. Hope MD¹⁵ | Ayse T. Kendi MD¹⁶ |
Samantha M. Thomas MS¹⁷ | Sherlonda Flem BS, CTR¹⁸ | James Brierley MB¹⁹ |
Elliot A. Asare MD, MS²⁰  | Kay Washington MD, PhD²¹ | Chanjuan Shi MD, PhD²² 

Summary of Three Notable Changes

- 1) Recommend **AGAINST** the use of **chromogranin A** for diagnosis and for treatment response
- 2) Incorporation of WHO grading of GEP-NET in staging, prognosis, and treatment decisions
- 3) Evolution of Stage 1 disease and role for endoscopic resection for NET of the stomach, duodenum, and colorectum

Chauhan A et al. Ca Cancer J Clin 2024; 74: 359-367


Treatments for GEP-NET

Types of Treatments for GI NET



Surgery

Catheter-based liver directed therapy




Habib A et al. Nat Rev Clin Oncol 2015.

Peptide Receptor Radionuclide Therapy (PRRT)/Radioligand Therapy (RLT)

Medical (Systemic) Therapies

- Targeted agents
- Chemotherapy options



Beck MN et al. Cancers 2022.

- Surgery preferred if can remove all disease; others mainly control disease
- Surgery can still be used even if disease can't be all removed (e.g. debulking liver surgery, resecting the primary tumor, etc)
- Treatments may be used separately, but sometimes can be used to compliment each other
- Each person's situation is different, and a multidisciplinary discussion is important

Localized GI NET: Primary Organ Site Influences Management

NET of small intestine (jejunum, ileum, colon)

- Bowel resection + lymph node resection

Pancreatic NET

- Pancreaticoduodenectomy (Whipple's)
- Observation (for smaller, stable tumors)

Gastric NET

Type 1 (~70-80%)

- Occurs with autoimmune atrophic gastritis (stomach parietal cells can't make acid)
- **Low** acid in stomach -> **high gastrin** -> growth of neuroendocrine cells
- Causes small, benign polyps (low chance of spread)
- **Monitor with endoscopies (i.e. no surgery or medical treatment)**

Type 2 (~5-10%)

- Occurs in context of Zollinger-Ellison Syndrome
- Tumor makes **high gastrin** -> **high acid**
- Medical treatment for acid secretion + resection tumor/stomach

Type 3 (~5-10%, Sporadic)

- Not Type 1 or Type 2
- Single, large tumor of the stomach with high chance for spread
- **Normal gastrin**
- Surgery + lymph node removal

Type 4

- Chronic proton pump inhibitors (PPI's)/antacids cause a similar physiology to Type 1
- **Low** acid in stomach -> **high gastrin** -> growth of neuroendocrine cells
- Often indolent and can be monitored with endoscopies


Appendiceal

- Simple appendectomy (<2 cm tumor)
- Right colon removal (hemicolectomy) + Lymph nodes (>2 cm)
- Debate over 1-2 cm tumors


Rectum

- Endoscopic resection if small tumors (up to 2 cm, if possible)
- Full rectal surgery + lymph nodes for larger tumors and/or abnormal lymph nodes on scans/biopsies

Therapies in GEP-NET



Systemic Therapies for GEP-NET



Dong M, Phan AT, Yao JC.
Clin Cancer Res, 2012; 18(7): 1830-6.

Pancreatic NET (pNET)

- Somatostatin analogs (SSAs) (octreotide LAR, lanreotide)
- PRRT/¹⁷⁷Lutetium (¹⁷⁷Lu)-dotatate
- Everolimus
- Capecitabine/Temozolomide (CAPTEM)
- Sunitinib
- Belzutifan (germline *VHL* only)
- *Teloristat ethyl (for refractory carcinoid syndrome diarrhea)

Other GI NET


- SSAs (octreotide LAR, lanreotide)
- PRRT/¹⁷⁷Lutetium (¹⁷⁷Lu)-dotatate
- Everolimus
- *Teloristat ethyl (for refractory carcinoid syndrome diarrhea)

- For symptom and tumor control
- Observation may be appropriate for patients with low grade/indolent disease
- **Based on current data, there is no established sequence of therapies**

Summary of Clinical Outcomes of GEP-NET Systemic Therapies


Objective response rate (ORR)	2-3%	<ul style="list-style-type: none"> pNET: 5 Other GI NET: 2% 	9.3%	<ul style="list-style-type: none"> pNET: 40% Other GI NET: ? (~4-50%) 	20-40% (↑ for higher grade disease and pNET)	18-50% (↑ in pNET)	pNET: 19% Non-pNET: 5%
Progression Free Survival (PFS, months)	Rate of 2 yr-PFS: 50-65%	<ul style="list-style-type: none"> pNET: 11.4 Other GI NET: 11-14 	11.4	<ul style="list-style-type: none"> pNET: 22.7 Other GI NET: ? (>30) 	>30	>30	pNET: 13.4 Non-pNET: 8.4
Overall Survival (OS, months)	Median not reached (NR)	Median NR	Median NR	pNET: 58.7 Other GI NET: ? (>20-30)	20-40	50-60	pNET: 40 Non-pNET: 22
Common Toxicities	Diarrhea, abdominal discomfort, gallstones	Mouth sores, diarrhea, fatigue, rash, hyperglycemia, potential pneumonitis	Diarrhea, nausea, fatigue, hand-foot syndrome	Cytopenias, nausea, diarrhea, fatigue, potential risk of MDS with long term temozolomide use	Cytopenias, nausea, fatigue, peripheral neuropathy	Nausea, fatigue, cytopenias, ~1.5% risk MDS/AML	Diarrhea, hypertension, hand-foot syndrome, blood clots

Systemic Therapy Considerations for GEP-NET



- No one size fits all with regards to treatment approach
- Multidisciplinary discussion important


Peptide receptor radionuclide therapy (PRRT): Revolutionized Treatment in GEP-NET




Becx MN et al. Cancers 2022; 14; 5792.

NETTER-1:

Improved PFS in Previously Treated WD-Gr 1/2 Midgut NET



No. at Risk	0	3	6	9	12	15	18	21	24	27	30
$^{177}\text{Lu-DOTATATE}$ group	116	97	76	59	42	28	19	12	3	2	0
Control group	113	80	47	28	17	10	4	3	1	0	0




Number at risk (number censored)	0	12	24	36	48	60	72	84	96
$^{177}\text{Lu-Dotatate}$ group	117 (0)	98 (9)	79 (12)	63 (13)	48 (16)	35 (17)	25 (22)	10 (35)	0 (44)
Control group	114 (0)	84 (8)	61 (14)	45 (18)	33 (23)	25 (26)	21 (27)	6 (39)	0 (45)

- ORR 18% vs 3%
- No OS benefit at final analysis, but 36% of pts on controlled arm received ^{177}Lu -dotatate at time of progression
- Median OS of 48.0 months in the ^{177}Lu -Dotatate group and 36.3 months (25.9–51.7) in the control group (HR 0.84 [95% CI 0.60–1.17]; two-sided $p=0.30$).
- Led to FDA approval in Feb 2018 for GEP-NET progressed on somatostatin analogs
- Data on non-midgut NET and higher grade disease lacking

NETTER-2: Improved PFS in 1L WD-Gr 2/3 GEP-NET


NETTER-2 (NCT03972488) is the first randomized trial to evaluate RLT as 1L treatment in any solid tumor



*Q8W during 177Lu-DOTATATE treatment then Q4W. †Octreotide LAR in retreatment phase is at discretion of investigator. ††, first line, G, grade, GEP-NET, gastroenteropancreatic neuroendocrine tumor, LAR, long-acting repeatable, Q8W, every 8 weeks, QOL, quality of life, R, randomization, RLT, radioligand therapy, SSTR, somatostatin receptor.

ASCO Gastrointestinal Cancers Symposium #GI24 PRESENTED BY: Simon Singh, MD, MPH

ASCO AMERICAN SOCIETY OF CLINICAL ONCOLOGY KNOWLEDGE CONQUERS CANCER



- First study demonstrating benefit of 177Lu-dotatate in 1L treatment of GEP-NET, including WD-Gr2/3 NET and for pancreatic NET
- ORR 43% vs 9.3%, with higher ORR in pancreas vs small bowel NET (51.2% vs 26.7%)
- Most common AE's are low grade diarrhea, abdominal pain, nausea
- ~1% risk of MDS/AML, which can occur within first 2 years
- Questions remain:
 - 1) Do all patients need 1L Lutathera?
 - 2) Is high dose octreotide LAR a fair comparator in higher grade disease?


Sequencing of Radioligand Therapy in GEP-NET

OCCULORANDOM STUDY

Academic randomized phase II trial in advanced progressive PanNET (well differentiated)



Inclusion between Feb 2015 – July 2020 in 10 French expert centers (GTE-RENATEN)



PARIS 2022 ESMO congress Eric Baudin


Content of this presentation is copyright and responsibility of the author. Permission is required for re-use.

Baudin E et al. Abstract 8870. ESMO Congress 2022

COMPETE STUDY

COMPETE trial design

Prospective, randomised, controlled, open-label, multi-centre phase 3 trial




Capdevila J E et al. ENETS 2025

Sequencing of Radioligand Therapy (RLT) in GEP-NET

OCCULORANDOM STUDY

Progression-free survival : real time blinded central review RECIST 1.1




PARIS 2022 ESMO congress Eric Baudin Content of this presentation is copyright of the author. Permission is required for re-use.

Baudin E et al. Abstract 8870. ESMO Congress 2022

COMPETE STUDY

PFS (primary endpoint by BICR)




Median PFS was significantly longer with n.c.a. ¹⁷⁷Lu-edotreotide vs everolimus (23.9 vs 14.1 months; p=0.022; HR 0.67, 95% CI [0.48, 0.95])

BICR, Blinded Independent Central Review; CI, confidence interval; HR, hazard ratio; n.c.a., non-carrier-added; PFS, progression-free survival

Capdevila J E et al. ENETS 2025

- OCULORANDOM and COMPETE suggest that earlier treatment with RLT is better (at least compared to sunitinib for pNET and everolimus for GEP-NET)
- Ongoing studies will address RLT compared to other systemic treatments (e.g. CAPTEM/FOLFOX) in higher-grade GEP-NET


Alpha-emitter PRRT: Evolving treatment in GEP-NET



Eur J Nucl Med Mol Imaging (2014) 41:2106–2119
DOI 10.1007/s00259-014-2857-9

Targeted Alpha-Emitter Therapy With ^{212}Pb -DOTAMTATE for the Treatment of Metastatic SSTR-Expressing Neuroendocrine Tumors: First-in-Human, Dose-Escalation Clinical Trial


Ebrahim S Delpassand, Izabela Tworowska, Rouzbeh Esfandiari, Julien Torgue, Jason Hurt, Afshin Shafie and Rodolfo Núñez
Journal of Nuclear Medicine January 2022. jnumed.121.263230, DOI: <https://doi.org/10.2967/jnumed.121.263230>



- Current PRRT treatments consists of beta-radiation
- Alpha-radiation particles penetrate cell with less scatter and increase chance for double stranded DNA breaks
- Preliminary studies suggest active even in patients who have received prior beta-emitter PRRT
- Phase 3 ACTION-1 Study ongoing against other systemic treatments (NCT05477576)


Tyrosine Kinase Inhibitors/Angiogenesis Inhibition in GEP-NET

- Tyrosine kinase inhibitors (TKI) target various pathways, particularly with respect to angiogenesis
- VEGF/VEGFR pathway is primary target but most TKI's target other pathways (e.g. c-KIT, PDGFR, others)
- New TKI's inhibit other pathways (e.g. FGFR, CSF1-R, MET, AXL) which may be able to overcome resistance after progression on prior treatments
- Sunitinib only approved TKI (for pNET)




Grillo F et al. Endocrine-Related Cancer, 2018; 25: R405–R418

Studies of Other Tyrosine Kinase Inhibitors




PRESENTED AT: Gastrointestinal Cancers Symposium | SIOGA is the property of the author, permission required for reuse. | PRESENTED BY: Garcia Carbonero, R, MD | #4021




PRESENTED AT: Gastrointestinal Cancers Symposium | SIOGA is the property of the author, permission required for reuse. | PRESENTED BY: Garcia Carbonero, R, MD | #4021

Garcia-Carbonero R et al. Abstract 360, ASCO GI 2019




PRESENTED AT: 2019 ASCO ANNUAL MEETING | #ASCO19 | PRESENTED BY: Emily Bergsland, MD | Abstract #4005




PRESENTED AT: 2019 ASCO ANNUAL MEETING | #ASCO19 | PRESENTED BY: Emily Bergsland, MD | Abstract #4005

Bergsland EK et al. Abstract 4005, ASCO 2019

- Various TKI's (pazopanib, axitinib, lenvatinib, surafatinib) have been evaluated
- New TKI's target other pathways (e.g. CSF-1R with surafatinib)
- No new approvals due to mixed results (pazopanib, axitinib), non-randomized & toxicities (lenvatinib), or need to be studied in other countries (surafatinib)



Capdevila J et al. J Clin Oncol 2021;39: 2304-2312.




Xiu J et al. Lancet Oncol 2020; 21: 1500-12.

Paulson S et al. Abstract 4114, ASCO 2021.

Alliance A021602/CABINET Study: Cabozantinib for Refractory GEP-NET and Pulmonary NET

CABINET Trial Study Design



Stratification factors:

- epNET: Concurrent SSA & Primary site (midgut GI/unknown vs. non-midgut GI/lung/other)
- pNET: Concurrent SSA & Prior sunitinib


Study Endpoints:

- **Primary Endpoint per cohort:**
 - Progression-free survival (PFS) by blinded independent central review
- **Secondary Endpoint per cohort:**
 - Overall survival
 - Objective response rate
 - Safety and tolerability




CABINET Trial Results


epNET Cohort: Progression-Free Survival (Local Review)




pNET Cohort: Progression-Free Survival (Local Review)



Best Overall Response (Local Review)



Overall Survival



Summary of CABINET Study Results

- Cabozantinib demonstrated significant improvement in PFS in previously treated GEP-NET
- Benefit seen in pancreas and small bowel NET, but benefit greater in pancreatic NET
- First study demonstrating benefit of an angiogenesis inhibitor in WD-Gr3 disease
- Notable AE's: Fatigue, diarrhea, hypertension, hand/foot syndrome
- Will be a new standard of care in previously treated GEP-NET (anticipated PDUFA date Apr 2025)
- Questions remain:
 - Optimal dose?: Recommended to start 60 mg po daily then dose-reduce as needed for toxicities
 - How best to sequence with other therapies? (e.g. pancreatic NET where there are more options)

Belzutifan in VHL-Associated Pancreatic NET (pNET)

The NEW ENGLAND JOURNAL of MEDICINE

ORIGINAL ARTICLE



Belzutifan for Renal Cell Carcinoma in von Hippel–Lindau Disease

	RCC (n=61)	Pancreatic Lesions (n=61)	pNET (n=22)	CNS Hemangioblastoma (n=50)
Objective Response Rate (ORR = CR + PR)	30 (49%)	47 (77%)	20 (91%)	15 (30%)
Complete response (CR)	0	6 (9.8%)	3 (13.6%)	3 (6.0%)
Partial response (PR)	30 (49%)	41 (67.2%)	17 (77.3%)	12 (24.0%)
Stable Disease (SD)	30 (49%)	12 (21.3%)	2 (9.1%)	31 (62.0%)
Median time to response	8.2 months	8.4 months	5.5 months	3.2 months
Median duration of response	Not reached	Not reached	Not reached	Not reached

- Belzutifan targets HIF-2^α which is upregulated in VHL associated cancers
- Secondary analysis of belzutifan in patients with VHL associated RCC who had concurrent pNET
- Belzutifan demonstrated ORR of pNET
- Listed in NCCN guidelines as an option for progressive VHL-associated pNET
- Remaining questions:
 - Most VHL pNET are indolent and localized disease --> when would Belzutifan be helpful?
 - Would this provide benefit in metastatic pNET (study excluded metastatic disease)?
 - Efficacy compared to other therapies for pNET? (i.e. sequencing)

Treatments for GEP-NEC

NEC Prevalence (SEER Database 1973-2012)



Dasari A et al. Cancer 2018

Current Treatment Paradigm in NEC

- Extrapolated from small cell lung cancer (SCLC) with use of platinum (cisplatin or carboplatin)/etoposide
- Data from retrospective series, except for 1 recent **prospective study** presented

Study	N	Histology (%)	Ki-67 Proportion	OS	PFS	RR
NORDIC-NEC ¹ (GI)	305	Small Cell: 38% Non-small cell: 49% Unknown: 13%	≥55%: 54%	11 mo	4 mo	Overall: 31% Ki-67 ≤ 55%: 15% Ki-67 ≥55%: 42%
FFCD-GTE ² (GI & unknown primary)	Total: 253 GI-NEC: 189	Small Cell: 39% Large Cell: 61%	51-80%: 47% >80%: 18%	11.6 mo	6.2 mo	50%
Mackey JR et al. ³ (GU)	Total 180 (106 bladder, 60 prostate, 8 renal, 6 ureter)	42.7% with mixed histology (adeno+ small cell);	Not reported	Overall: 10.5 mo Prostate: 7 mo Bladder: 13 mo	?	?
Margolis B et al. ⁴ (Cervix)	1,896	Not reported	Not reported	~10 mo	?	?
Morizane C et al. (GI, prospective)⁵	170	Small Cell: 48% Large Cell: 52%	Ki-67 ≥50%: 85%	12.5 mo	5.6 mo	54.5%

¹Sorbye H et al. Ann Oncol 2013 ²Walter T et al. Eur J Cancer 2017 ³Mackey J et al. J Urol 1998 ⁴Margolis B et al. Gynecol Oncol 2016 ⁵Morizane C et al. JAMA Oncol 2022.

Systemic Therapies for Refractory NEC

NEC
FOFLOX
FOLFIRI
FOLFIRINOX
Temozolomide +/- Capecitabine
Nivolumab + Ipilimumab
Pembrolizumab (if MSI-H or TMB ≥ 10)


- No established standard for NEC after progression on platinum/etoposide
- Outcomes remain poor:
 - ORR: 20-30%
 - Median PFS ~3 months
 - Median OS 6-9 months
- Need more effective treatments

Adapted from NCCN Neuroendocrine Guidelines, v. Feb 2022

Nivolumab + Ipilimumab in Refractory Extrapulmonary NEC

Response Rate by Tumor Grade of Neuroendocrine Neoplasm


SWOG S1609: DART Study



Patel SP et al. Clin Cancer Res 2020; 26: 2290-6

GCO-001 NIPINEC trial

Randomized, non-comparative phase II trial – Fleming's two-stage design



Walter T et al. Abstract LBA41. ESMO Congress 2021


- Effective in high grade NEN/NEC (and not NET)
- Extrapulmonary ORR: 10-25% (lower compared to lung NEC ~20-60%; nivolumab alone ORR: 7.1%)
- 6-month PFS: 20-30%; Median OS 6-11 months
- **Suggests that combination therapy is more effective than monotherapy**

SWOG S2012: Randomized Phase 2/3 Trial of First Line Platinum/Etoposide +/- Atezolizumab for Extrapulmonary NEC

- **Activated Dec 2, 2021** (restricted to small cell NEC)
- **Jan 2023: Amended to allow all NEC Subtypes**
- **Feb 2024: Amended to remove Ki-67 for GU NEC**

Key Eligibility: (N= 118/189)

- Metastatic poorly-differentiated extrapulmonary (i.e. exclude lung) NEC with Ki-67 ≥ 55%
- Evaluable, measurable and non-measurable disease
- Zubrod PS 0-2
- No prior treatment **EXCEPT one cycle of platinum/etoposide allowed**
- Asymptomatic brain metastases eligible
- **Stratification factors:**
 - 1) PS 0-1 vs 2
 - 2) Known prostate vs GI vs other origin




Primary endpoint: OS (from time of randomization)

Secondary endpoints: OS (from time of maintenance/observation), PFS, ORR, DOR


Translational analyses: Banking tissue and blood for future biomarker analyses

Novel Therapies for NEC


Bi-specific T-cell engagers (BiTE)¹



Antibody Drug Conjugates (ADC)²



CAR-T³



¹Einsele H et al. Cancer 2020.


²Dumontet C et al. Nature Rev Drugs Disc. 2023; 22; 641-661

³Du B et al. Cancer Cell 2025; 43; 665-679


SEZ6 is highly expressed in neuroendocrine neoplasms (NENs) and is the target of the ADC ABBV-706

- SEZ6 is a neuroendocrine lineage marker specifically expressed in SCLC and high-grade NENs¹
- ABBV-706 is a first-in-class ADC composed of a potent Top1i attached via a stable linker to a SEZ6-directed monoclonal antibody, delivering the cytotoxic payload specifically to SEZ6-expressing tumor cells
- Preliminary data of ABBV-706 monotherapy showed manageable safety and promising activity in SCLC and NENs²

SEZ6 Gene Expression



ABBV-706 Mechanism of Action




1. Wiedemeyer WR, et al. *Mol Cancer Ther.* 2022;21:986-998. 2. Chandana S, et al. ASCO 2024. Abstract 3001 (oral presentation).

ADC, antibody-drug conjugate; LCNEC, large cell NEC; NEC, neuroendocrine carcinoma; NEPC, neuroendocrine prostate cancer; NET, neuroendocrine tumor; SCLC, small cell lung cancer; SEZ6, seizure-related homolog protein 6; Top1i, topoisomerase 1 inhibitor.

ABBV-706 has promising antitumor activity in NENs

Change in Target Lesion Size by Dose (N=65)



Outcome	Total NEN (N=65) ^{a,b}
ORR, ^c n (%) [95% exact CI]	24 (36.9) [25.3, 49.8]
Best response, n (%)	
CR	1 (1.5)
PR	23 (35.4)
SD	37 (56.9)
PD	4 (6.2)
Median DOR, months [95% CI]	6.37 [4.44, 9.46]
Median PFS, months [95% CI]	7.62 [5.52, 8.31]

- As of January 3, 2025, cutoff, 17/65 (26%) patients remain on treatment
- Median follow-up was 8.67 months for the 2.5-mg/kg cohort (n=49/65, 75%)

^aPatients with response evaluation per RECIST v1.1. ^b49 of 65 (75%) patients received ABBV-706 at 2.5 mg/kg. ^cRequires a CR or PR confirmed ≥ 4 weeks later. CI, confidence interval; CR, complete response; DOR, duration of response; NEN, neuroendocrine neoplasm; ORR, objective response rate; PD, progressive disease; PFS, progression-free survival; PR, partial response; RECIST v1.1, Response Evaluation Criteria in Solid Tumors version 1.1; SD, stable disease.



Thank you

dbzhen@uw.edu